

Immune Globulin Autoimmune Disorder Referral Form

Field Contact:

PATIENT INFORMATION

| | | |
|---|--|--|
| Patient Name: _____ | Phone: (____) - ____ - _____ | Emerg. Contact: _____ |
| Date of Birth: ____/____/____ <input type="checkbox"/> Male <input type="checkbox"/> Female | Email: _____ | Emerg. Phone: (____) - ____ - _____ |
| SSN: ____ - ____ - ____ | Preferred method of contact: <input type="checkbox"/> Phone <input type="checkbox"/> Email | |
| Physical Address: _____ | Height: _____ in Weight: _____ lb | Date: ____/____/____ |
| City: _____ State: _____ Zip: _____ | Allergies: _____ | Medications: _____ (Please attach additional pages if necessary) |

PRESCRIPTION BENEFITS INFORMATION (Please attach front and back of insurance card)

| | | | | |
|------------------|------------|----------------|--------------|--------------|
| Plan name: _____ | ID#: _____ | Group #: _____ | RxBIN: _____ | RxPCN: _____ |
|------------------|------------|----------------|--------------|--------------|

PRESCRIBER INFORMATION

| | | |
|-------------------------------------|------------------------------|--|
| Prescriber Name: _____ | Phone: (____) - ____ - _____ | PREVIOUS THERAPY Medication(s): _____ _____ <input type="checkbox"/> IV <input type="checkbox"/> SC Rate: _____ |
| Address: _____ | Fax: (____) - ____ - _____ | |
| City: _____ State: _____ Zip: _____ | License #: _____ | |
| Contact: _____ | NPI #: _____ | |
| Clinic/Hospital Affiliation: _____ | Medicaid Provider #: _____ | |

CLINICAL INFORMATION - PRIMARY DIAGNOSIS - ICD-10

| | |
|--|---|
| <input type="checkbox"/> Acute Infective Polyneuritis (Guillain-Barre Syndrome) ICD-10 _____ | <input type="checkbox"/> Myasthenia Gravis without (Acute) Exacerbation ICD-10 _____ |
| <input type="checkbox"/> Chronic Inflammatory Demyelinating Polyneuropathy (CIDP) ICD-10 _____ | <input type="checkbox"/> Pemphigus (Pemphigus Foliaceus, Pemphigus Vulgaris) ICD-10 _____ |
| <input type="checkbox"/> Dermatomyositis ICD-10 _____ | <input type="checkbox"/> Pemphigoid ICD-10 _____ |
| <input type="checkbox"/> Inflammatory Polyneuropathy, Unspecified ICD-10 _____ | <input type="checkbox"/> Polymyositis ICD-10 _____ |
| <input type="checkbox"/> Multiple Sclerosis (MS) ICD-10 _____ | <input type="checkbox"/> Stiff-Person Syndrome ICD-10 _____ |
| <input type="checkbox"/> Multifocal Neuropathy (MMN) ICD-10 _____ | <input type="checkbox"/> Other: _____ ICD-10 _____ |
| <input type="checkbox"/> Myasthenia Gravis with (Acute) Exacerbation ICD-10 _____ | |

PRESCRIPTION AND ORDERS

Administer: ☐ SCIG ☐ IVIG Product: ☐ Pharmacist to determine (or) ☐ Formulation: _____

Dose: (please select option(s) and provide complete information, pharmacy to round the nearest 5 gram vial)

- ☐ Loading Dose: _____ gm/kg OVER _____ day(s), then ☐ Maintenance Dose: _____ gm/kg OVER _____ day(s), EVERY _____ week(s) x _____ Refills
- ☐ Other Regimen: _____

Infusion Rate: (please select one and provide complete information)

- ☐ Pharmacist to determine
- ☐ Start at _____ mL/hr, then increase by _____ mL/hr every _____ minutes to maximum rate _____ mL/hr

Access: ☐ Peripheral ☐ PICC ☐ Port ☐ Other: _____

IV Maintenance (Flushing): _____ Dispense Quantity Sufficient with 11 Refills - or - _____ Decline

- Sodium Chloride 0.9% 10mL Prefilled Syringe: Flush IV access device with sodium chloride 3-10mL to maintain line patency.
- Heparin 10 units/mL 5mL Prefilled Syringe: Flush peripheral IV access device with Heparin 10 units/mL 1-5mL as needed to maintain line patency. (pt<10kg)
- Heparin 100 units/mL 5mL Prefilled Syringe: Flush peripheral IV access device with Heparin 100 units/mL 3-5mL as needed to maintain line patency. (pt>10kg)

Pre-Treatment: _____ Dispense Quantity Sufficient with 11 Refills - or - _____ Decline

- Acetaminophen 325mg Tablet: 1-2 tablets by mouth 15-30 minutes before each infusion.
- Diphenhydramine 25mg Capsule: 1-2 capsules by mouth 15-30 minutes before each infusion.
- ☐ Other: _____

Adverse/Anaphylactic Reactions: Anaphylaxis kit to be used in the event of anaphylactic reaction per HPC Protocol and will contain the following:

- Diphenhydramine 25mg Capsule #4
- Diphenhydramine 50mg/mL 1mL vial #2
- Sodium Chloride 0.9% 500mL Bag #1
- Epinephrine Injection Auto-Injector 0.3mg (>30kg pt) or 0.15mg (<30kg pt) Two-Pack #1
- Sodium Chloride 0.9% 10mL Prefilled Syringe #4

Ancillary Supplies: Dispense ancillary supplies and equipment (including pump and pole) needed to provide home infusion therapy.

Nursing Orders for Home Infusion (IV Only)

Frequency of vital signs monitoring:

a. If initial infusion or more than 8 weeks since last infusion:

- Prior to infusion
- q 15 minutes for the first hour
- q 30 minutes during the second hour
- q 1 hour for the remainder of the infusion

b. If subsequent infusion within 8 weeks:

- Prior to infusion
- q 15 minutes for the first hour
- q 1 hour for the remainder of the infusion

PRESCRIBER SIGNATURE (Stamp signature not allowed, physician signature required)

Product Selection Permitted

Dispense as Written

Date



PHONE: 1-800-757-9192
www.hpcinfusion.com

PLEASE INCLUDE ALL MEDICAL RECORDS & LAB VALUES
PLEASE FAX TO 1-855-813-0583

All rights in the product names of all third-party products appearing in this document, whether or not appearing with the trademark symbol, belong exclusively to their respective owners.

IGAD_0109-1021